

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 33/569, C12Q 1/04, G01N 33/543	A1	(11) International Publication Number: WO 96/38729 (43) International Publication Date: 5 December 1996 (05.12.96)
(21) International Application Number: PCT/SE96/00721 (22) International Filing Date: 31 May 1996 (31.05.96) (30) Priority Data: 9502024-4 2 June 1995 (02.06.95) SE (71) Applicant (for all designated States except US): PHARMACIA BIOSENSOR AB [SE/SE]; S-751 82 Uppsala (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): WAHLSTRÖM, Lennart [SE/SE]; Ekhagsvägen 8, S-743 40 Storvreta (SE). PATEL, Pradip [GB/GB]; 207A Firtree Road, Epsom Downs, Surrey KT17 3LB (GB). HAINES, John [GB/GB]; 33 Strathcona Avenue, Bookham, Surrey KT23 4HW (GB). (74) Agents: WIDÉN, Björn et al.; Pharmacia AB, Patent Dept., S- 751 82 Uppsala (SE).		(81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: PATHOGEN ASSAY METHOD (57) Abstract A method for detecting a pathogen in a sample comprises the steps of: (i) optionally treating the sample to obtain a pathogen-enriched sample, (ii) mixing a predetermined amount of the optionally pathogen-enriched sample with a solution containing a predetermined amount of antibody capable of specifically binding to the pathogen to permit the antibody to bind to the pathogen, (iii) separating any pathogen from the mixture to obtain a pathogen-free solution, and (iv) determining the amount of antibody present in the pathogen-free solution to thereby detect binding of antibody to the pathogen as indicative of the presence of pathogen in the sample.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

PATHOGEN ASSAY METHOD

The present invention relates to a novel assay method for detecting pathogens, especially in foods.

5 The statistics of food poisoning incidences in recent years, the strict food safety legislation and introduction of quality systems in food manufacturing has increased the need for real-time techniques for pathogen analysis during food production.

10 The conventional cultural method for the detection of pathogens like *Salmonella* and *Listeria* requires at least five days to isolate and identify the organism from a food product. In foods, *Salmonella* and *Listeria* are often present in low numbers, largely outnumbered by competing
15 organisms, and occasionally in a stressed state. The method therefore includes an initial non-selective pre-enrichment stage designed to allow the resuscitation and growth of the
~~target pathogen. The subsequent selective enrichment stage~~
utilises growth media that are inhibitory to competitor
20 cells whilst still maintaining or allowing the growth of the target organism. The selective enrichment process aims to improve the ratio of *Salmonella* or *Listeria* to competitor cells. Colonies are then isolated from the enriched culture by plating on to selective and
25 differential agar media. Any presumptive-positive colony is then confirmed biochemically and identified serologically. While cultural techniques for the microbiological examination of foods have high sensitivity (capable of detecting one viable cell), they are time-consuming and
30 highly labour- and material-intensive.

Considerable research has therefore been focused on the development of rapid methods for the detection of pathogens in foods. However, none of the rapid methods available commercially so far, such as ELISA, impedance,
35 and DNA hybridisation, give results in real-time but all require at least 24 to 48 h of cultural enrichment in order to increase the productivity of the pathogens above the

detection limits of these techniques, which at present is above 10^5 cfu/ml.

The available immunoassays of the enzyme-linked immunosorbent assay (ELISA) type for *Salmonella*, for example, are to be performed on selectively enriched samples (48-h culture) and comprise heat-solubilising *Salmonella* antigen, adding the antigen-containing broth to antibody-coated wells of a microtitre plate and incubating, adding antibody-enzyme conjugate, and after incubation adding substrate and reading absorbance.

The object of the present invention is to provide a pathogen assay which together with a rapid assay protocol has a sufficiently low detection limit, such as 10^2 - 10^4 cfu/ml, to permit detection of pathogens in contaminated foods without 24-h selective enrichment of the food sample, and therefore within a single working day (less than 24 h).

In accordance with the present invention it has now been found that this object may be achieved by a modified inhibition type immunoassay wherein a possibly pathogen-containing sample suspension is first reacted with a predetermined excess level of antibodies against the pathogen. The suspension is then subjected to a separation procedure to remove the pathogen, and the resulting antibody level in the separated solution is determined to thereby indirectly detect the pathogen. A detected low relative level of antibody will indicate a high pathogen level in the reacted sample, whereas a detected high level of antibody will indicate a low pathogen level in the sample.

Accordingly, the present invention in a broad aspect provides a method for detecting a pathogen in a sample, comprising the steps of:

- (i) optionally treating the sample to obtain a pathogen-enriched sample (in case of pathogen in the sample),
- (ii) mixing a predetermined amount of the optionally pathogen-enriched sample with a solution containing a predetermined amount of an antibody capable of specifically

binding to the pathogen to permit the antibody to bind to the pathogen,

(iii) separating any pathogen from the mixture to obtain a pathogen-free solution, and

5 (iv) determining the amount of antibody present in the pathogen-free solution to thereby detect binding of antibody to the pathogen as indicative of the presence of pathogen in the sample.

The antibody is preferably added in excess of the
10 amount corresponding to the (expected) maximum amount of pathogen to leave unbound antibody in the reaction mixture.

The term pathogen as used herein comprises not only bacteria, but also virus, fungi and protozoa.

Although the detection of foodborne pathogens is an
15 important application of the present invention, the method may, of course, also be applied to samples of other origin, such as, e.g., samples for clinical assays, e.g., blood, urine, etc.

Exemplary of common foodborne pathogens are *Salmonella*
20 and *Listeria*. Adequate pre-enrichment (and optionally 6-h selective enrichment) procedures for food samples to reach a pathogen level of about 10^2 - 10^4 cfu/ml as mentioned above are well-known to those skilled in the art.

The removal of the pathogen from the antibody reaction
25 mixture is preferably performed by filtration or centrifugation. Suitable filters for this purpose are known to a person skilled in the art and will not be described in detail herein.

The technique chosen for detecting the excess antibody
30 in the pathogen-free solution is not critical per se. In a preferred detection technique, however, the solution is contacted with an optical sensor surface having immobilized thereon a receptor, such as an antibody, for the primary antibody, and the change in refractive index at the surface
35 related to binding of primary antibody present in the solution to the sensor surface is measured.

Advantageously, the contacting of the sample with the surface is performed by passing the sample over the surface utilizing a liquid flow system, i.e. a flow cell.

If desired, further sensitivity in the assay may be
5 obtained by using a secondary reagent capable of binding to the primary antibody bound to the sensor surface. Optionally, also a tertiary reagent may be used which binds to the secondary reagent.

Binding of the receptor, such as an antibody, to the
10 surface may be carried out in conventional ways well-known to those skilled in the art. If, for example, the optical surface has a polymeric organic layer at its surface, the receptor may be directly covalently bound to the surface using known linker reagents. Alternatively, an intermediate
15 ligand, such as an antibody, which binds the receptor may first be covalently bound to the surface before this bound intermediate ligand is exposed to the receptor for the
~~analyte to bind this to the surface. It will be appreciated~~

~~that covalent and/or affinity bonding may be effective in~~
20 binding the receptor to the surface. However, when the receptor is bound to the surface, it is important that its ability to bind the ligand, i.e. in this case the primary antibody, should remain unchanged.

The term antibody as used herein is to be interpreted
25 broadly. Thus, in addition to a whole antibody, the antibody may be a fragment thereof, such as an Fab fragment, an Fv fragment, a single chain fragment (scFv), a single heavy chain or even a peptide (based on the nucleotide sequence of the antibody gene) having binding
30 activity. The antibodies which may be used in the invention may be obtained by conventional methods and are many times commercially available. Although polyclonal antibodies may conveniently be used in the method of the invention, monoclonal antibodies may be preferred at least in certain
35 cases for their greater specificity.

The measurement of the change in refractive index at the surface may advantageously be based on evanescent wave sensing, such as surface plasmon resonance spectroscopy

(SPRS), Brewster angle refractometry, critical angle refractometry, frustrated total reflection (FTR), evanescent wave ellipsometry, scattered total internal reflection (STIR), optical wave guide sensors, evanescent wave based imaging, such as critical angle resolved imaging, Brewster angle resolved imaging, SPR angle resolved imaging, etc. In the currently preferred method for carrying out the invention, the measurement is based on surface plasmon resonance. This technique is described, inter alia, in EP-A-0305109, EP-A-0267142 and WO-A-90/05295. The optical surface which is used in the measurement based on surface plasmon resonance preferably comprises a gold film and a hydrogel bound to the gold film, as described in WO 90/05303. This type of optical surface may easily be regenerated so that a single surface may be used for many analyses. The overall cost per analysis can therefore be reduced considerably. Suitable apparatus incorporating such an optical surface is the BIAcore® system available from Pharmacia Biosensor AB, (Uppsala, Sweden) the methods of operation of which are described in the BIAcore® Methods Manual (Pharmacia Biosensor AB). In the BIAcore® system, a flow system passes the sample over a replaceable sensor chip forming one wall of a flow cell. The sensor chip supports a gold layer which typically has a thickness of 50 nm. A carboxylated dextran is bound to the gold layer via a linker layer. To this dextran layer the receptor for the primary antibody may be bound.

The invention will now be described in more detail with reference to the following Examples and the accompanying drawings, wherein:

Fig. 1 is a diagram showing the dose-response curve for *S. enteritidis* and the response of *C. freundii* obtained by the method of the invention in the same biosensor flow cell;

Fig. 2 is diagram showing the dose-response curve for *S. typhimurium* and the response of *C. freundii* obtained by

the method of the invention in the same biosensor flow cell;

Fig. 3 is a diagram showing the dose-response curve for *S. napoli* and the response of 5 non-salmonellae obtained by the method of the invention in the same biosensor flow cell;

Fig. 4 is a diagram showing the dose-response curve for *S. stanley* and the response of 3 non-salmonellae obtained by the method of the invention in the same biosensor flow cell;

Fig. 5 is a diagram showing the dose-response curve for *S. thompson* and the response of 4 non-salmonellae obtained by the method of the invention in the same biosensor flow cell;

Fig. 6 is a diagram showing the dose-response curve for *S. typhimurium* and the response of *S. enteritidis*, *C. freundii* and *E. coli* obtained by the method of the invention in the same biosensor flow cell and using a monoclonal as primary antibody;

Fig. 7 is a diagram showing the dose-response curves for three *Listeria* serotypes and the response of three non-listeriae obtained by the method of the invention; and

Fig. 8 is a diagram showing the dose-response curves for five *Salmonella* serotypes and the response of 10 non-salmonellae obtained using a commercial prior art *Salmonella* ELISA (*Salmonella*-Tek; Organon).

The analyses in the Examples were carried out on a BIAcore® system (Pharmacia Biosensor AB, Uppsala, Sweden) with a Sensor Chip CM5 as the optical sensor surface.

EXAMPLES

MATERIALS AND METHODS

Buffers and regeneration solutions

Hepes buffered saline (HBS, Pharmacia Biosensor AB), containing 10 mM Hepes, pH 7.4, 150 mM sodium chloride (NaCl), 3.4 mM ethylenediaminetetraacetic acid (EDTA), and 0.005% v/v Surfactant P20, was used as the standard running buffer in all experiments.

Sodium acetate buffer (10 mM, pH 5) was used as the coupling buffer to enable immobilisation of the antibodies to the sensor surface.

The following solutions were tested for their ability to regenerate the sensor surface: formic acid (1M), hydrochloric acid (10-100 mM HCl), glycine/HCl (10 mM glycine titrated with HCl to required pH), ethanolamine/HCl (1M ethanolamine titrated with HCl to required pH), potassium hydroxide (10-100 mM KOH), potassium chloride (4 M KCl), sodium hydroxide (10-100 mM NaOH), KCl/NaOH (4 M KCl titrated with NaOH to required pH), glycine/NaOH (10 mM glycine titrated with NaOH to required pH), ethanolamine/NaOH (1 M ethanolamine titrated with NaOH to required pH), and sodium dodecyl sulphate (0.05-1 % SDS).

15 Antibodies

Bactrace anti-Salmonella CSA-1 (Kirkegaard & Perry Laboratories), below referred to as "KPL Bactrace"

~~Mouse anti-Salmonella broad specificity Mab (Serotec)~~

Bactrace anti-Listeria (genus specific, Dynatech)

20 Rabbit anti-goat IgG_{fab} (RAG_{Fab}, Pierce)

Goat anti-mouse IgG_{fab} (GAM_{Fab}, Pierce)

Immobilisation of antibodies to the sensor surface

Pharmacia Biosensor's recommended protocol was used for this purpose and involved the reagents (NHS, EDC and ethanolamine) contained in the amine coupling kit (code no. 25 BR-1000-50, Pharmacia Biosensor AB).

The carboxylated dextran surface was initially activated using a 35 µl (flow rate of 5 µl/min) injection of N-hydroxysuccinimide (NHS) and N-ethyl-N¹(dimethylaminepropyl)carbodiimide (EDC) in distilled water (0.05 M NHS/0.2 M EDC), across a flow cell surface. This was then followed by an injection (35 µl) of the antibody in 10 mM sodium acetate buffer, pH 5, and a subsequent injection (35 µl) of 1 M ethanolamine 35 hydrochloride.

Microorganisms

The following organisms were used: *Salmonella enteritidis* P167808, *S. typhimurium* NCTC 74, *S. napolii* NCTC

6853, *S. stanley* R23, *S. thompson* R24, *Listeria innocua* GP100/100, *L. monocytogenes* NCTC 5105, *L. innocua* LFRA isolate B047, *L. seeligeri* LFRA B052, *Bacillus cereus* NCTC 11145, *Staphylococcus aureus* NCTC 4136, *Staph. aureus* LFRA isolate B3, *Staph. epidermidis* LFRA isolate PB14, *Yersinia enterocolitica* NCTC 11174, *Escherichia coli* NCTC 9001, *E. coli* NCTC 9112, *E. coli* NCTC 8545, *Proteus mirabilis* 1102/11/1, *Citrobacter freundii* NCTC 6266 and *Micrococcus luteus* NCTC 7495.

10 All the cultures were grown at 37°C for 24 h in Trypticase Soya Broth (TSB). The cultures were then diluted (serial decimal) in HBS and enumerated by plating 0.1 ml on to Trypticase Soya Agar (TSA).

Regeneration of the sensor surface

15 The regeneration conditions were optimised for each anti-*Salmonella* and anti-*Listeria* antibody tested. This was achieved by screening a range of potential regeneration solutions, as specified in the BIAcore™ user manual, in order to find the best possible solution that completely dissociated the antibody from the sensor surface. Thus, the capture antibody, e.g. RAGFab, was immobilised to the sensor surface and the anti-*Salmonella* antibody, e.g. KPL Bactrace, was injected using the manual command mode to allow binding to the biosensor surface. Then a weak regeneration solution was injected for 1 min and the degree of regeneration monitored. It was found that for KPL Bactrace, for example, the best regeneration solution was 50 mM NaOH.

EXAMPLE 1: Inhibition assay for *Salmonella*

30 Determination of minimum relative response

Varying levels of an anti-*Salmonella* antibody were reacted with a high level (approximately 10^8 cfu/ml) of a *Salmonella* culture in order to determine the level of primary antibody required to give a minimum relative response in the BIAcore®.

Preparation of bacterial cultures

Bacterial cultures were grown at 37°C for 24 h in trypticase soya broth (TSB). The 24-h cultures were then

heated in boiling water for 20 min. Prior to assay, the heat-inactivated cultures were diluted in Hepes Buffered Saline (HBS).

Optical biosensor assay

5 500 μ l of diluted primary antibody to *Salmonella*, corresponding to the above predetermined level, were pre-reacted with 500 μ l of heat-inactivated bacterial cells and mixed for 15 min on a shaker (Luckham multimix). The mixture was then filtered with a 0.22 μ m Sartorius filter
10 to remove excess primary antibody from antibody bound to the bacterial cells, and 15 μ l of the filtrate containing the separated excess primary antibody were injected over the sensor surface coated with the corresponding capture antibody. The sensor surface was then regenerated. The time
15 to first result was 30 min. The higher the level of *Salmonella*, the lower was the relative response, and vice versa. The results are shown in Figs. 1 to 6.

 Figs. 1 to 5 show obtained response curves for *S. enteritidis*, *S. typhimurium*, *S. napoli*, *S. stanley* and *S.*
20 *thompson*, respectively, together with the response of various non-salmonellae challenged at approximately 10^8 cfu/ml in the same flow cell. In this case, a polyclonal antibody, KPL Bactrace, was used as primary antibody. Rabbit anti-goat IgG_{fab} fragment (RAG_{fab}) was used as
25 capture antibody on the sensor surface, and regeneration was performed with 50 mM sodium hydroxide, 2x1 min pulses. As appears from the Figs. 1-5, the detection limit for all five *Salmonella* serotypes was in the range 10^2 - 10^4 cfu/ml.

 Fig. 6 shows the detection of *Salmonella* using a
30 monoclonal anti-*Salmonella* (Serotec) as primary antibody and goat anti-mouse IgG_{fab} fragment (GAM_{fab}) as capture antibody on the sensor surface. Regeneration of the GAM_{fab} sensor surface reacted with the primary antibody was performed using a combination of 1x15 μ l formic acid (1M),
35 1x5 μ l NaOH (2.5 mM) and 1x15 μ l formic acid (1M) pulses. It is seen from the dose-response curve for *S. typhimurium* shown in Fig. 6 that no cross-reaction was observed for *C. freundii* and *E. coli* tested at approximately 10^8 cfu/ml. A

S. enteritidis control at approximately 10^8 cfu/ml was also successfully detected.

EXAMPLE 2: Inhibition assay for *Listeria*

In the same way as described for *Salmonella* above, an inhibition assay for three *Listeria* strains was performed using a RAGFab-coated sensor surface and a polyclonal anti-*Listeria* as primary antibody. The results are shown in Fig. 7. As appears therefrom, the sensitivity level of the assay was approximately 10^2 - 10^4 cfu/ml, with no cross-reaction observed for *E. coli*, *Staph. aureus* and *Staph. epidermidis* tested at approximately 10^8 cfu/ml.

EXAMPLE 3: Commercial *Salmonella* ELISA (comparative)

Bacterial cultures were prepared as in Example 1 above, and five *Salmonella* serotypes and responses of ten non-*Salmonella* were detected by a commercial *Salmonella* assay, "Salmonella-Tek ELISA", using the following protocol.

100 μ l samples were pipetted into wells of a microtitre plate. The samples were then incubated at 37°C for 30 min. After washing six times with wash solution, 100 μ l conjugate were added and incubated at 37°C for 30 min. The wells were then washed six times, and 100 μ l of substrate were added and incubated at room temperature for 30 min. 100 μ l of stop solution were then added and the absorbance was read at 450 nm. The time to first result was 2 h.

The results are shown in Fig. 8, demonstrating a detection limit of approximately 10^4 - 10^6 cfu/ml for salmonellae, with considerable cross-reaction from *C. freundii* and *E. coli* R6, but not eight of the other non-salmonellae all tested at approximately 10^8 cfu/ml.

CLAIMS

1. A method for detecting a pathogen in a sample,
comprising the steps of:
 - 5 (i) optionally treating the sample to obtain a pathogen-enriched sample,
 - (ii) mixing a predetermined amount of the optionally pathogen-enriched sample with a solution containing a predetermined amount of antibody capable of specifically
10 binding to the pathogen to permit the antibody to bind to the pathogen,
 - (iii) separating any pathogen from the mixture to obtain a pathogen-free solution, and
 - (iv) determining the amount of antibody present in the
15 pathogen-free solution to thereby detect binding of antibody to the pathogen as indicative of the presence of pathogen in the sample.

2. The method according to claim 1, wherein the antibody
20 is added in step (ii) in a sufficient amount to leave unbound antibody in the mixture.
3. The method according to claim 1 or 2, wherein the determination of antibody in step (iv) comprises contacting
25 at least a predetermined portion of the pathogen-free solution with an optical sensor surface having immobilized thereon a receptor for the antibody, and measuring a change in refractive index at the surface related to the amount of antibody present in the pathogen-free solution contacted
30 with the sensor surface.
4. The method according to claim 1, 2 or 3, wherein the separation of the pathogen in step (iii) is performed by
filtration.
- 35 5. The method according to any one of claims 1 to 4, wherein said receptor is an antibody.

6. The method according to any one of claims 1 to 5, wherein the pathogen is a foodborne pathogen and step (i) comprises enriching the pathogen in the food sample to a level in the range of from about 10^2 to about 10^4 cfu/ml.
- 5 7. The method according to any one of claims 1 to 6, wherein said measurement of the change of refractive index is based on internal reflection.
- 10 8. The method according to claim 7, wherein said measurement of the change of refractive index is based on surface plasmon resonance.
- 15 9. The method according to claim 8, wherein said optical sensor comprises a gold film and a hydrogel bound to the gold film.
-
- 20 ~~10. The method according to any one of claims 1 to 9,~~
wherein the sample is contacted with the optical sensor surface by passing the sample over the sensor surface in a liquid flow.
- 25 11. The method according to any one of claims 1 to 10, wherein the pathogen is selected from *Salmonella* and *Listeria*.

1/8




FIG. 1

2/8




FIG. 2

3/8




FIG. 3

4/8




FIG. 4

5/8




FIG. 5

6/8




FIG. 6

7/8




FIG. 7

8/8





FIG. 8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 96/00721

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: G01N 33/569, C12Q 1/04, G01N 33/543

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: G01N, C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

BIOSIS, MEDLINE, DBA, WPI, EDOC, PAJ, SCISEARCH, PATENT CITATION INDEX

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Journal of Food Protection, Volume 53, No 10, October 1990, Céline Morissette et al, "Simple and Rapid Inhibition Enzyme Immunoassay for the Detection of Staphylococcal Enterotoxin B in Foods" page 834 - page 840	1-2,4,6
A		3,5,7-11
A	International Journal of Food Microbiology, Volume 12, 1991, S. Notermans et al, "Immunological methods for detection of foodborne pathogens and their toxins" page 91 - page 102	1-11
A	EP 0496345 A1 (NAGASE & COMPANY, LTD.), 29 July 1992 (29.07.92)	1-11

☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

- * Special categories of cited documents:
- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed
- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
28 August 1996	11 -09- 1996
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86	Authorized officer Patrik Andersson Telephone No. +46 8 782 25 00

Information on patent family members

05/09/96

International application No.

PCT/SE 96/00721

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A1- 0496345	29/07/92	CA-A- 2059690 JP-A- 5005744	23/07/92 14/01/93
<hr/>			